


# Greenhouse Gases, Climate Change, Carbon Sequestration, and Carbon Credit Trading




#### Areas to Cover:


- Characteristics of Light
- Earth's Radiation Balance
- Greenhouse Effect
- Greenhouse Gases
- Agricultural Mitigation of GHG's
- Methods to Estimate GHG's



To understand why greenhouse gases in the atmosphere are an issue, we must first discuss radiation transfer within the atmosphere.



Solar energy is made up partly of visible light (ROYGBIV), as well as some ultraviolet and infrared light.



Some radiation from the sun is reflected by the earth, the rest is absorbed. This absorbed radiation heats the earth surface, which then in turn emits heat/thermal radiation back to the atmosphere and space.






# What happens to light at the earth's surface?

- Reflected (albedo)
- Absorbed
  - Heat Surface
  - Heat/Evaporate Water
  - Photosynthesis





- Shortwave Radiation
  - From Sun
  - Visible Light
  - Reflected and Absorbed by Earth
- Longwave Radiation
  - Emitted from Earth
  - Heat (Thermal)




The simple equation for net radiation balance: Rn - Net Radiation, SW is shortwave (solar) radiation, LW is longwave (terrestrial) radiation. The greenhouse effect provides the incoming longwave radiation.





#### Greenhouse Effect and Greenhouse Gases

The **greenhouse effect** is the process in which the absorption and emission of thermal infrared radiation (heat) by a planet's atmosphere warms that planet's surface.








Earth's average temperature is


15 C (59°F)





Without the greenhouse effect, the Earth's average temperature would be 0°F




The big concern with increasing atmospheric concentrations of greenhouse gases is that the greenhouse gas absorption portion of the radiation balance with increase, trapping more heat in the atmosphere and increasing the global temperature, thus global warming.

















#### What are Some Greenhouse Gases?

- CO<sub>2</sub>
- H<sub>2</sub>O
- N<sub>2</sub>O
- Ozone
- CH<sub>4</sub>
- Fluoromethane compounds




Carbon Cycle. Graphic shown to depict the wide array of sources and sinks of carbon in the environment. Note that while soils are an important storage point for carbon, the oceans, in particular the deep oceans, have the largest overall carbon storage of any mechanism.



Graphic representing the nitrogen cycle. Identifies the nitrification/denitrification reactions as the primary source of N2O.



- For accounting purposes, greenhouse gases are counted in carbon dioxide equivalents:
  - $CO_2 = 1 CO_2 equivalent$
  - CH<sub>4</sub> ~ 20 CO<sub>2</sub> equivalents
  - N<sub>2</sub>O ~ 300 CO<sub>2</sub> equivalents










### Agricultural Sources of GHG's

- CO<sub>2</sub>
  N<sub>2</sub>O
  CH<sub>4</sub>



Anthropogenic agricultural sources of carbon dioxide. This list does not include natural sources of CO2 like soil respiration, microbial activity, natural decomposition processes.







#### Offsetting GHG Emissions

- Emissions Reductions (CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O)
- Carbon Sequestration (CO<sub>2</sub>)





#### **Emissions Reductions**

- Fewer, or combined trips/passes
- Reducing tillage
- Alternate/renewable fuels





### Reduce CO<sub>2</sub> Emissions

- Reduced Trips
  - Reduced Tillage
  - No-Till
  - Combining Operations





N2O emissions can be reduced by splitting applications of nitrogen fertilizers, applying only for crop needs, and utilizing nitrification inhibitors. Storing manures dry can help reduce N2O losses from solid manure piles.






## Methane Reduction and Capture

- Use an aerobic manure system
- Capture methane with digester technology
  - Emits carbon dioxide, and generates energy










#### **Predictive Tools**

- CCX recommendations
- IPCC recommendations
- 1605(b) techniques
- Process models (Century, EPIC)
- Web-based (COMET-VR, COLE)



Map of zones for carbon sequestration rates from CCX.




#### Cropland (not certified in OR)

#### Allowed (follow NRCS-329):


- · no-till drill
- no-till and strip-till planters
- low disturbance liquid manure injectors, anhydrous ammonia applicator, manure knife applicator

#### Not Allowed:

- field cultivators, tandem disk, offset disk, chisel plow, moldboard plow
- Burning or residue removal



For recently planted grasses (after 1999).








## Forestry

- Based on stand type/age
- > stems per acre




http://illinoisclimate.org/calculator.php



# Intergovernmental Panel on Climate Change (IPCC)

- CO<sub>2</sub> emissions factors based on land use, tillage, and inputs
- $N_2O = .016 * N applied$
- Applicable for national accounting




http://www.cometvr.colostate.edu



#### **COMET-VR**

- Version 1.1
- 226 MLRA's
- 20-40 rotation/cropping choices per MLRA
- 12 soil textures
- Century model w/ improved uncertainty estimate





Required Responses to use COMET-VR: The tool requires 7 inputs by the user based on information that we feel land managers only know. The location is needed to determine climate data for the model run. Parcel information is for the user to identify which lands they are working on. Soils information is based on soil texture and hydric class which allows the model to capture artificially drained areas. We are only dealing with mineral soils in the version of COMET-VR. The management history is selected by the user and is limited at this time, but is currently being expanded to capture the systems as reported in the National Resources Inventory. Once all the data is entered, the user selects Get Carbon and the tool delivers a response in 5 seconds or less.





## Required Responses to Utilize COMET-VR


- Management History (crop rotations, tillage systems or grazing systems)
  - Pre 1970's
  - 1970's-1990's
  - Base: 1990's-Current
  - Reporting Period: Current + 10 years































## **Carbon Credit Trading**

- Certified emissions reductions/sequestration
  - measured/modeled
- Aggregator Entity
- Trading Mechanism



#### Aggregators

- · Iowa Farm Bureau
- Farmers Union (North Dakota)
- · Illinois Conservation and Climate Init.
- Beartooth Capital Partners (Montana)
- The Carbon Fund
- Environmental Credit Corporation
- Other new ones every day



## Trading – Chicago Climate Exch.

 "Chicago Climate Exchange (CCX) is North America's only and the world's first global marketplace for integrating voluntary legally binding emissions reductions with emissions trading and offsets for all six greenhouse gases."



Example of a carbon trade. Farmer no-tills, gains carbon credits. Sells them to an aggregator. The aggregator markets the credits on the Chicago Climate Exchange. Offsetters can purchase credits on CCX to market as greenhouse gas offsets. Private travelers can purchase credits to offset the "carbon footprint" of their travel.



#### Review:

- Sunlight and Thermal Radiation
- Greenhouse Effect and Global Temperature
- GHG's and Agriculture
- Methods of Estimating GHG Emission Reductions and Sequestration
- Carbon Credit Trading

